The Effect of Sulfur on the Color of Triphenyl Methane Dyes and Certain Phthaleins

STANFORD UNIVERSITY LIBRARIES
WITHDRAWN

DISSERTATION

SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF THE JOHNS HOPKINS UNIVERSITY IN CONFORMITY WITH REQUIREMENTS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY

BY

HAROLD STEPHENS HOLT

781
65H

BALTIMORE
February, 1924

The Effect of Sulfur on the Color of Triphenyl Methane Dyes and Certain Phthaleins

---

DISSERTATION

SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF
THE JOHNS HOPKINS UNIVERSITY IN CONFORMITY
WITH REQUIREMENTS FOR THE DEGREE
OF DOCTOR OF PHILOSOPHY

BY

HAROLD STEPHENS HOLT

BALTIMORE
February, 1924

ACKNOWLEDGMENT

Acknowledgment is due to Dr. E. Emmet Reid at whose suggestion this work was undertaken and under whose guidance it was carried out. His friendly advice and valuable suggestions call for the most cordial thanks.

Acknowledgment is also due to Drs. Frazer, Patrick, Lovelace and Thornton for instruction received in the classroom and laboratory.

PART I

The Effect of Sulfur on the Color of Triphenyl Methane Dyes

THE EFFECT OF SULFUR ON THE COLOR OF TRIPHENYLMETHANE DYES

Introductory and Historical

Within the last several years, a considerable amount of work has been done in this Laboratory on the auxochrome effect of alkyl-linked sulfur, more particularly in the field of azo dyes.[1]

The present investigation is a further extension of this idea into the field of triphenylmethane dyes. A search through the literature showed few triphenylmethane dyes containing alkyl-linked sulfur.

Brand and Stallmann[2] have studied the auxochrome effect of alkyl mercapto groups on the phenol ethers of triphenylmethane. They have prepared compounds of the type $(C_6H_5)_2C(OH)C_6H_4SCH_3$ (*o*) and $C_6H_5C(HO)(C_6H_4SCH_3)_2$ (*o*) and have shown that such compounds in acetic acid or alcoholic solution when treated with strong sulfuric or hydrochloric acid, develop color. These cannot be classed as true dyes although they do show the auxochrome effect of the $—SCH_3$ group.

[1] (a) Waldron and Reid, *J. Am. Chem. Soc.*, **45**, 2399 (1923). (b) Foster and Reid, *ibid.*, **46**, 1936 (1924).

[2] Brand and Stallmann, *Ber.*, **54**, 1578 (1921).

The Present Investigation

The Grignard reagents were prepared from the alkyl ethers of *o*- and *p*-bromo-thiophenol and made to react with Michler's ketone:[3] $RS.C_6H_4MgBr + [(CH_3)_2NC_6H_4]_2CO \longrightarrow [(CH_3)_2NC_6H_4]_2C(OH)C_6H_4.SR$. The desired *o*- and *p*-bromo-thiophenol ethers were obtained from the corresponding amine compounds by the Sandmeyer reaction.

The dyes so made are derived from malachite green by the substitution of —SR in the *ortho* or *para* position in the third ring. The groups used were methyl, *iso*propyl and *iso*amyl in the *para* position and methyl in the *ortho*. Dyes containing the phenyl and benzyl sulfide or methylsulfone groups could not be prepared, since the reaction between magnesium and the bromophenyl-thio-ethers appears to be confined to the simpler members of the series for neither *p*-bromophenyl-benzyl sulfide nor *p*-bromophenyl methylsulfone could be made to attack the magnesium. In the case of *p*-bromophenyl-thiophenol a reaction did take place after long heating but a brown solid immediately precipitated which did not react with Michler's ketone. Insufficient material was available to enable us to study this reaction further.

For comparison, dyes were made containing the methyl and methoxy groups in the same positions by the action of the Grignard reagents from *o*- and *p*-tolyl iodide and *o*- and *p*-iodo-anisole.

Wool skeins were dyed in the usual way using 1% of the dye, 3% of sulfuric acid and 10% of Glauber's salt. The colors[4] are given in Table I.

TABLE I

COLORS OF SUBSTITUTED MALACHITE GREENS ON WOOL

Substituent	Ortho	Para
None	Yellowish-green	Yellowish-green
$-CH_3$	Dull bluish-green (weak)	Bright yellowish-green
$-OCH_3$	Dull yellowish-green (weak)	Dull yellowish-green
$-SCH_3$	Bluish-green (turquoise)	Dull reddish-blue
$-SCH(CH_3)_2$	.....	Dull blue-gray (weak)
$-SC_5H_{11}$(*iso*)	.....	Dull yellowish-green (weak)

The introduction of sulfur in the *para* position has a pronounced deepening effect on the color and the shift is toward the blue, that is, the shorter wave length. This effect is especially noticeable with the lower members of the series.

As the molecular weight of the alkyl group attached to the sulfur atom increases, a peculiar slate gray results, making it difficult to determine the color. Certainly some of the auxochrome effect is lost.

[3] Baeyer and Villiger, *Ber.*, **36**, 2789 (1903). Ehrlich and Sachs, *Ber.*, **46**, 4296 (1903). Votocek and Matejzka, *Ber.*, **46**, 1756 (1913).

[4] Thanks are due Dr. R. E. Rose of E. I. du Pont de Nemours and Co. for the color designations.

In the *ortho* position all of the substituents diminish the intensity of the dye and all shift the color towards the blue, the —SCH_3 group showing the greatest effect.

Experimental Part

Preparation of *p*-Nitro-thiophenol.—The procedure of Waldron and Reid was modified slightly, as more consistent results were obtained and less time was required. It was found to be unnecessary to acidify the reaction mixture and then redissolve the free mercaptan. The procedure was as follows.

One gram molecular weight of *p*-nitro-chlorobenzene (157 g.) was treated with sodium disulfide as described by Waldron and Reid. The reaction mixture was poured into about 300 cc. of water and the mixture filtered at once into saturated sodium hydroxide solution, precipitating most of the sodium salt of the mercaptan and some disulfide. This mixture was filtered through cloth and the mercaptan removed by treating with warm water containing a little sodium sulfide. When saturated sodium hydroxide is added to this solution the product is precipitated quite pure.

The disulfide may be collected and reduced with sodium sulfide in strongly alkaline solution (for 125 g. of disulfide, 35 g. of sodium sulfide and 35 g. of sodium hydroxide in 150 cc. of water are required). This reaction mixture is treated exactly as above. The success of the preparation depends on the length of time the reaction mixture is allowed to boil before pouring into water and this is best determined by a few trials. The yields were about 60% on the first step.

Ethers of *o*- and *p*-Bromo-thiophenol, $RS.C_6H_4Br$.—The ethers of *o*- and *p*-amino-thiophenol were made according to the methods of Waldron and Reid and of Foster and Reid. The amino groups in these were replaced by bromine as follows.

A solution of 25 g. of the hydrochloride of the aminophenyl thio-ether in about 300 cc. of water containing 40 g. of concd. hydrochloric acid was cooled to 0° and diazotized in the usual manner. The diazo solution was then poured into a suspension of 22 g. of cuprous bromide and 25 g. of potassium bromide in a little water at 80°. After evolution of nitrogen had ceased, 100 cc. of concd. hydrochloric acid was added and the mixture steam distilled. The distillate was made strongly alkaline to remove any phenol and the bromo compound extracted with benzene. It was distilled in a vacuum after it had been dried and the solvent evaporated. As the phenyl and benzyl ethers were not volatile with steam the reaction mixture, after diazotizing and decomposing as before was extracted with benzene. The benzene solution was shaken with 10% alkali, separated, dried and the solvent evaporated. This left the bromo compounds mixed with considerable tarry material. From this mixture they were distilled under reduced pressure and further purified by redistilling.

***p*-Bromophenyl-Methylsulfone.**—A solution of 22 g. of *p*-aminophenyl methylsulfone, in 300 cc. of water containing 40 g. of concd. hydrochloric acid was diazotized in the usual manner. The diazo solution was poured into a suspension of 20 g. of cuprous bromide and 25 g. of potassium bromide in 50 cc. of water at 80°. This solution was heated to boiling and filtered hot. The bromosulfone crystallizes from the filtrate in yellow needles. By extracting the residue on the filter with hot water and evaporating to small volume, more product can be obtained.

The yields, properties and analyses of the new compounds prepared are

given in Table II. Analyses for sulfur were made with the Parr bomb and densities were determined with the Ostwald pycnometer.

TABLE II

PROPERTIES AND ANALYSES OF *o*- AND *p*-BROMOPHENYL THIO-ETHERS, $RS.C_6H_4Br$

Group	Position	B. p. or m. p. °C.	Density 25/25	Sulfur Calcd. %	Sulfur Found %	Yield %
—$SCH(CH_3)_2$	*o*	b. p., 130–135 (11 mm.)	1.2804	13.85	13.59	35
—SC_6H_5	*o*	b. p., 175–177 (12 mm.)	1.3733	12.09	12.17	50
—SCH_3	*p*	m. p., 27	....	15.76	15.71	55
—$SCH(CH_3)_2$	*p*	b. p., 120 (11 mm.)	1.2338	13.85	13.62	60
—$SCH_2CH(CH_3)_2$	*p*	b. p., 140–143 (15 mm.)	1.1467	12.35	12.23	37
—$SCH_2C_6H_5$	*p*	m. p., 48	....	11.47	11.38	50
—SO_2CH_3	*p*	m. p., 97.5	....	13.61	13.79	50

Preparation of the Dyes

The following procedure was used in all cases. About 10 g. of the bromo thio-ether in absolute ether was treated with magnesium turnings and a crystal of iodine. The magnesium dissolved slowly in all cases and it was necessary to heat the mixture at 35° overnight. The calculated quantity of Michler's ketone in dry benzene was then added, precipitating an orange-colored solid which turned green on contact with air. This mass was separated from the solvent by filtration or decantation and dissolved in 1:1 hydrochloric acid. After the mixture had been filtered, saturated zinc chloride solution was added and the dye precipitated by the addition of solid sodium sulfate.

The reaction with *o*-bromophenyl-thiophenol.—To 8.2 g. of this compound and 0.75 g. of magnesium turnings was added 15 cc. of ether and a crystal of iodine. The mixture was heated at 35° overnight. Solution of the magnesium suddenly took place after 24 hours and a brown solid precipitated. More ether was added, followed by the calculated amount of Michler's ketone in dry benzene. The precipitate which formed did not turn green on exposure to air, and on dissolving in hydrochloric acid only colorless products were formed. No dye could be obtained.

Method of Dyeing

One-tenth g. of dye was dissolved in about 500 cc. of warm water containing 1 g. of sodium sulfate and 0.3 g. of sulfuric acid. After solution of the dye, any insoluble material was removed by filtration. A 10 g. skein previously soaked in hot water was then immersed and the solution gradually brought to boiling, the total time of immersion being one-half hour in each case. The skein was then removed, washed with water and air-dried.

Summary

1. A number of new bromophenyl thio-ethers have been prepared. On coupling with Michler's ketone these gave dyes in which the auxochrome effect of the alkyl mercapto groups could be observed.

2. Sulfur in the *para* position has a decided auxochrome effect on the color and the shift is toward the blue. Sulfur in the *ortho* position has a similar effect but to a less degree. Increasing the molecular weight of the alkyl groups results in a loss of some of the auxochrome effect. In general the auxochrome effect of the various groups is in the order $—SCH_3 > —OCH_3 > —CH_3$.

PART II

The Effect of Sulfur on the Color of Certain Phthaleins

THE EFFECT OF SULFUR ON THE COLOR OF CERTAIN PHTHALEINS

Introductory and Historical

It has been shown[1] that the introduction of sulfur into triphenylmethane dyes causes a decided shift in color, this change being toward the shorter wave length. This idea has been extended into the field of phthaleins, the problem in general being the synthesis of a series of phthaleins containing alkyl-linked sulfur and observation of the auxochrome effect.

No mention of phthaleins containing alkyl sulfur could be found in the literature although compounds such as phenolsulfonephthalein and its derivatives have been known for some time.[2]

The Present Investigation

The method of attack first suggesting itself was the fusion of phthalic anhydride with phenols containing sulfur substituents, $RS.C_6H_4.OH$, the procedure being similar to that used in the preparation of phenolphthalein. These phenols were made from the corresponding amines[3] by the diazo reaction: $RS.C_6H_4.NH_2 \longrightarrow RS.C_6H_4.OH$.

It was found, however, that when one mole of phthalic anhydride is fused with two moles of *o*-hydroxyphenyl methyl sulfide at 180° in the presence of fused zinc chloride, hydroxymethyl-mercapto-anthraquinone is formed. That is, although two moles of thio-ether were used only one reacted: $C_6H_4(CO_2)O + RS.C_6H_4.OH \longrightarrow C_6H_4(CO)_2C_6H_2(OH)SR$. No attempt was made to determine the position of the hydroxyl or methyl mercapto groups.

o-Hydroxyphenyl-thiophenol, $HO.C_6H_4.SC_6H_5$, does not react with phthalic anhydride under similar conditions.

[1] *J. Am. Chem. Soc.*, **46**, 2329 (1924).
[2] Remsen, *Am. Chem. J.*, **20**, 263 (1898).
[3] Foster, *J. Am. Chem. Soc.*, **46**, 1936 (1924).

Orndorff and Murray[4] and Moir[5] have shown that *p*-hydroxybenzoyl-*o*-benzoic acid has the power of condensing with a great variety of phenols to give mixed phthaleins. It has been found that this acid will condense with such compounds as thiophenol, *o*-nitro-thiophenol, *p*-nitro-thio-anisole, *o*-nitro-thio-anisole and the *o*-hydroxyphenyl thio-ethers. The present work was confined almost entirely to the hydroxy thio-ethers and by this reaction it is possible to prepare phthaleins containing alkyl-linked sulfur.



The procedure consisted in heating *p*-hydroxybenzoyl-*o*-benzoic acid with an excess of the hydroxy thio-ether in the presence of fused zinc chloride. The products so formed are true phthaleins; they form acetates, benzoates and oximes and in alkaline solution are deeply colored. Neither the acetates nor benzoates could be crystallized from the common solvents. Analyses were made on the oximes which could be obtained quite pure.

Table I contains the new hydroxyphenyl thio-ethers which were prepared and fused with *p*-hydroxybenzoyl-*o*-benzoic acid, their properties and analyses.

TABLE I

HYDROXYPHENYL THIO-ETHERS AND METHYLSULFONE

Compound	B. p. or m. p. °C.	Density 25/25	Sulfur Calcd.	Sulfur Found	Yield %
o-HO.$C_6H_4SCH_3$	b. p., 105 (22 mm.)	1.168	22.85	22.60	52
o-HO.$C_6H_4SC_4H_9$	b. p., 125 (10 mm.)	1.061	17.58	17.73	45
o-HO.$C_6H_4SC_6H_5$	m. p., 44	...	15.84	15.53	50
o-HO.$C_6H_4.SO_2CH_3$	m. p., above 220	...	18.60	18.41	..

The indicators prepared are all derivatives of phenolphthalein containing the —SR or —SO_2R group adjacent to one of the hydroxyls. The indi-

TABLE II

SUBSTITUTED PHENOLPHTHALEINS, THEIR SÖRENSEN (*P*H) RANGE[a] AND COLOR IN ALKALI WITH MELTING POINTS AND ANALYSES OF THEIR OXIMES

Group	*P*H values	Color in alkali	M. p. °C.	Sulfur Calcd. %	Sulfur Found %
—SCH	8.4–10	violet	189 decomp.	8.44	8.54
—SC_4H_9	8.6– 9.8	violet	175–176 decomp.	7.60	7.56
—SC_6H_5	9.0–10	violet-pink	190 decomp.	7.25	6.98
—SO_2CH_3	9.6	orange	165–167 decomp.	7.78	7.50

[a] Thanks are due the LaMotte Chemical Products Co. for the determination of the *P*H values.

[4] Orndorff and Murray, *J. Am. Chem. Soc.*, **39**, 678 (1917).

[5] Moir, *Trans. So. African Acad.*, **1917–1922**.

cators could not be obtained in a condition pure enough for analysis, the sulfur determinations running 0.7 to 1.2% low, but they are readily converted into oximes. These are easily obtained pure by recrystallization and may be relied on for characterization. Analyses of the oximes are given.

Under similar conditions, *p*-hydroxybenzoyl-*o*-benzoic acid condenses with thiophenol, *o*-nitro-thiophenol, *p*-nitro-thio-anisole and *o*-nitro-thio-anisole. The products dissolved in alkali forming deep red or red-violet solutions but proved difficult to purify and were not further studied.

Discussion of Results

The effect of the alkyl mercapto groups on the color was observed by adding 1 cc. of a 0.02% indicator solution in alcohol to 20 cc. of a buffer solution of P_H 12. A similar solution of phenolphthalein was used for comparison.

In general the results agree with those obtained by us on triphenylmethane dyes.[1] The introduction of the— SCH_3 group results in a purple color, that is, a shift toward the blue. As the molecular weight of the alkyl group increases, much of the purple is lost until finally with the indicator containing the —S—C_6H_5 group, it has almost disappeared and a violet-pink results. Oxidation to the sulfone destroys whatever auxochrome effect the sulfur had, and the color is orange and of low tinctorial power. The usefulness of this indicator is diminished for this reason.

The alkyl mercapto groups have little effect on the P_H values of the indicators. An examination of the tables will show them all to have nearly the same range as phenolphthalein, turning a little further on the alkaline side.

Experimental Part

The Preparation of the *o*-Hydroxyphenyl Thio-ethers.—A solution of 25 g. of the corresponding amine hydrochloride in about 250 cc. of water containing 40 g. of concd. hydrochloric acid was cooled to zero degrees and diazotized with a slight excess over the calculated amount of sodium nitrite. The diazo solution was then warmed, the thio-ether distilled with steam and extracted from the distillate with benzene. The benzene solution was dried with sodium sulfate, the solvent evaporated and the thio-ether distilled under reduced pressure. In the case of the phenyl-thio-ether which is not volatile with steam, the reaction mixture was dried and the residue distilled in a vacuum.

***o*-Hydroxyphenyl Methylsulfone.**—Thirty-four g. of *o*-aminophenyl methylsulfone was diazotized and the diazo solution decomposed by boiling. The material came out of solution as a tarry mass which was separated and dissolved in sodium hydroxide solution. This was filtered and the filtrate diluted to about 800 cc. and acidified with acetic acid. The solid which precipitated could not be crystallized but was dissolved in alcohol and reprecipitated by the addition of ether.

p-Hydroxybenzoyl-*o*-benzoic acid was made by boiling phenolphthalein oxime with dil. sulfuric acid (1:8) as described by Orndorff and Murray.[4]

The Condensation of Phthalic Anhydride with *o*-Hydroxyphenyl Methyl Sulfide

A mixture of 5 g. of phthalic anhydride and 9.4 g. of thio-ether was heated to 180° for several hours in the presence of fused zinc chloride; it became viscous and solidified on cooling. The solid was powdered and boiled with dil. hydrochloric acid to remove basic zinc salts. This treatment converted the solid to a black, tarry substance which again solidified on cooling. It was dissolved in sodium hydroxide solution, the solution filtered and reprecipitated with strong acetic acid. It was a dark solid dissolving in alkalies forming a reddish-purple solution. Attempts to crystallize it from the usual solvents were unsuccessful. A partially purified product was obtained by dissolving in ether and filtering into petroleum ether in which it is insoluble. Under these conditions it was precipitated as a salmon-colored amorphous solid that melted at 188° with decomposition.

Anal. Calc. for $C_6H_4(CO)_2C_6H_2(OH)SCH_3$: S, 11.88. Found: 12.30.

The Condensation of *p*-Hydroxybenzoyl-*o*-benzoic Acid with *o*-Hydroxyphenyl Thio-ethers

The following procedure was used in the preparation of each indicator. About 10 g. of acid and a slight excess of thio-ether were heated to 180° in a porcelain or metal dish for several hours. A little fused zinc chloride was added to speed the reaction. The mixture became dark and viscous and solidified on cooling. After it had been pulverized it was dissolved in water, the solution made acid with hydrochloric acid and the excess of thio-ether removed by steam distillation. The solid material was filtered off, dissolved in 10% sodium hydroxide solution and this solution again filtered to remove any zinc salts. The alkaline filtrate was diluted to a considerable volume and made acid with acetic acid, precipitating the free indicator. The yields are practically quantitative.

Acetylation and Benzoylation

Acetates were prepared by boiling with acetic anhydride for three hours and pouring into water. The resulting products were pasty and could not be made to crystallize. The benzoyl derivatives were prepared according to the directions of Bistrzycki and Nencki[6] but could not be purified.

Conversion of the Indicators to their Oximes

The procedure followed was similar to that described by Orndorff and Murray. These oximes were pure enough for analysis.

Summary

1. A number of new indicators have been prepared by fusing *p*-hydroxybenzoyl-*o*-benzoic acid with *o*-hydroxyphenyl thio-ethers, and by comparison with phenolphthalein the auxochrome effect of the alkyl mercapto groups has been studied.

[6] Bistrzycki and Nencki, *Ber.*, **29**, 132 (1896).

2. Sulfur in the *ortho* position has a decided auxochrome effect, the shift being toward the blue. As the molecular weight of the alkyl group increases, much of this effect is lost. Oxidation of sulfide to sulfone destroys whatever auxochrome effect the sulfur had.

BIOGRAPHY

The writer, Harold Stephens Holt, was born January 17, 1897 at Fargo, North Dakota. He received his early education at the public schools there and in 1915 entered the State Agricultural College from which he received the degree of B.Sc. in 1919. The same year he entered Harvard University and received the degree of M.A. from that institution in 1921. During the year 1920–21 he was assistant in chemistry at Radcliffe College. He enrolled at the Johns Hopkins University as a graduate student in chemistry in the fall of 1921 and has been in attendance continuously since that time.

SEP 8 1926